

# **PyGAMMA:**

## **A Python Interface To GAMMA**

Jaiwen Chen<sup>†</sup>, Mikolai Fajer<sup>†</sup>, Peter Fajer<sup>†^</sup>, Khaled Khairy<sup>†\*</sup>, and Scott A. Smith<sup>†</sup>

<sup>\*</sup> Department of Chemistry, Northeastern University, Boston, Massachusetts

<sup>^</sup> Department of Molecular Biology, Florida State University, Tallahassee, Florida 32310

<sup>†</sup> National High Magnetic Field Laboratory, Florida State University, Tallahassee, Florida 32310

Running Title: PyGAMMA: A Python Interface to GAMMA

Corresponding Author: Scott A. Smith  
National High Magnetic Field Laboratory  
1800 E. Paul Dirac Drive  
Tallahassee, Floirda, 32310  
USA  
Tel: (850) 644 6348  
Fax: (850) 644 1368  
Email: sasmith@magnet.fsu.edu

To be submitted to J. Magn. Reson.

## ABSTRACT

A Python interface has been constructed for the GAMMA magnetic resonance simulation platform. Through utilization of the BOOST library and simple code additions we were able to export virtually the entire GAMMA package into Python. This effectively exports the full functionality of the platform into an interactive environment without sacrificing its object oriented nature or flexibility. Herein we discuss the implementation and its advantages as well as demonstrate its use with several examples.

*Key Words:* GAMMA; Python, BOOST, simulations; numerical calculations

## INTRODUCTION

The GAMMA magnetic simulation (MR) platform (12). Its versatility stems directly from its object oriented design. Users must have some knowledge and C++ programming and be able to Furthermore, since programs built using GAMMA are compiled into standalone executables, changes made to program variables may demand a time consuming recompilation before the user can see the effects of said changes.

## PYTHON AND GAMMA

## EXAMPLES

As a first step

[1]

Fig. 1. Calculat

Equation [1] satisfactorily explains why the minima for the different values of  $\alpha$  in Fig. 1 (a) to 40.7  $\mu\text{m}$  under the conditions employed, may not be a suitable measure of the thickness of the slice excited by a  $90^\circ$  pulse.

Corresponding simulations to those presented in Figures 3 and 4 were performed for sinc and cosine shaped pulses. Key features for all four pulse types are summarised in Table 1.

**TABLE 1**  
**Averaged features of the 2D plots of  $A_{\text{max}}$ ,  $A_{\Delta}$  and  $A_{\text{ratio}}$  for  $t_p \geq 10$  is for rectangular (Fig. 3), Gaussian (Fig. 4), sinc and cosine shaped pulses**

Feature	Pulse Shape			
	rectangular	Gaussian	sinc <sup>b</sup>	cosine

- 
- a* The errors given for  $A_{\max}$  and  $A_{\Delta}$  are based on calculations performed at  $1^\circ$  intervals. This was sufficient resolution to produce a variation in results for rectangular pulses, but not in results for the other pulse shapes studied.
- b* The results for sinc shaped pulses reflect the calculated echo shapes which have several local maxima and minima even at relatively small flip angles.

Experiments to test the calculated results were performed at  $\Omega/2\pi = 111.5$  MHz in an accurately calibrated STRAFI gradient of  $12.091 \pm 0.016$  T/m (13) using a probe that can take samples up to 5 cm in more conventional STRAFI experiment using  $f_b = 256,000$  Hz. The  $A_{\text{ratio}}$  values measured experimentally for each component of the phantom at specific  $t_p$  values close to 90 degrees, 120 degrees, 135 degrees and 180 degrees are summarised in Table 2.

## DISCUSSION

The calculated  $A_{\max}$  values in Figures 3 and 4 show that the maximum echo amplitude for rectangular and lobe of the sinc excitation envelope. It is evident that the sample outside of this range makes a significant contribution to echo amplitude.

## CONCLUSIONS

It is now possible to calibrate pulses for STRAFI echo trains accurately to within 1% in a simple. The GAMMA platform includes relaxation treatments and their use in STRAFI echo train calculations are now under investigation.

## ACKNOWLEDGEMENTS

GAMMA is provided to the public courtesy of its authors and is supported by the National High Magnetic Field Laboratory. We wish to express our gratitude to the REU exchange program for providing the means. Detailed information and downloading of PyGAMMA is available at <http://gamma.magnet.fsu.edu> .

## REFERENCES

3. B. Geil, Measurement of translational molecular diffusion using ultrahigh magnetic field gradient NMR, *Concepts in Magn. Reson.* **10**, 299-321 (1998).
12. S. A. Smith, T. O. Levante, B. H. Meier, and R. R. Ernst, Computer simulations in magnetic resonance. An object oriented programming approach, *J. Magn. Reson. A* **106**, 75-105 (1994).